

A Constrained Mixed Effects Model Based on Semilinear Differential Equation for Cell Polarity Signaling in Tip Growth of Pollen Tubes

Zhen Xiao

University of California, Riverside, USA, zxiao001@ucr.edu

Nicolas Brunel

Université d'Evry Val d'Essonne, FRANCE nicolas.brunel@ensiie.fr

Zhenbiao Yang

University of California, Riverside, USA zhenbiao.yang@ucr.edu

Xinping cui*

University of California, Riverside, USA xinping.cui@ucr.edu

The key of tip growth in eukaryotes is the polarized distribution on plasma membrane of a particle named ROPI. This distribution is the result of a positive feedback loop, whose mechanism can be described by a Differential Equation parametrized by two meaningful parameters k_{pf} and k_{nf} . In this paper, we introduce a mechanistic Integro-Differential Equation (IDE) derived from a spatiotemporal model of cell polarity and we show how this model can be fitted to real data i.e ROPI intensities measured on pollen tubes. At first, we provide an existence and uniqueness result for the solution of our IDE model under certain conditions. Quite interestingly, this analysis gives a tractable expression for the likelihood, and our approach can be seen as the estimation of a constrained nonlinear model. Moreover, we introduce a population variability by introducing a constrained nonlinear mixed model. We then proposed a constrained Least Squares method to fit the model under single subject case, and two methods, constrained Methods of Moments and constrained Restricted Maximum Likelihood (REML) to fit the model under the multiple subjects case. The performances of all the three methods are studied in a simulation example and are used on a real multiple subjects dataset

Keywords: Constrained Maximum Likelihood, Semilinear Differential Equation, Integro-Differential Equation, Mixed Models